arXiv:1604.04557v1 [math.NT] 15 Apr 2016

REVERSED DICKSON POLYNOMIALS OF THE FOURTH KIND OVER FINITE FIELDS

KAIMIN CHENG[A,B], SHAOFANG HONG[*,A] AND XIAOER QIN[C]
[A]MATHEMATICAL COLLEGE, SICHUAN UNIVERSITY, CHENGDU 610064, P.R. CHINA
[B]DEPARTMENT OF MATHEMATICS, SICHUAN UNIVERSITY JINJIANG COLLEGE, PENGSHAN 620860, P.R. CHINA
[C]SCHOOL OF MATHEMATICS AND STATISTICS, YANGTZE NORMAL UNIVERSITY, CHONGQING 408100, P.R. CHINA

Abstract. In this paper, we obtain several results on the permutational behavior of the reversed Dickson polynomial $D_{n,3}(1,x)$ of the fourth kind over the finite field $\mathbb{F}_q$. Particularly, we present the explicit evaluation of the first moment $\sum_{a\in\mathbb{F}_q} D_{n,3}(1,a)$.

1. Introduction

Let $\mathbb{F}_q$ be the finite field of characteristic p with q elements. Associated to any integer $n \geq 0$ and a parameter $a \in \mathbb{F}_q$, the n-th *Dickson polynomials of the first kind and of the second kind*, denoted by $D_n(x,a)$ and $E_n(x,a)$, are defined for $n \geq 1$ by

$$D_n(x,a) := \sum_{i=0}^{[\frac{n}{2}]} \frac{n}{n-i}\binom{n-i}{i}(-a)^i x^{n-2i}$$

and

$$E_n(x,a) := \sum_{i=0}^{[\frac{n}{2}]} \binom{n-i}{i}(-a)^i x^{n-2i},$$

respectively, and $D_0(x,a) := 2, E_0(x,a) := 1$, where $[\frac{n}{2}]$ means the largest integer no more than $\frac{n}{2}$. In 2012, Wang and Yucas [6] further defined the *n-th Dickson polynomial of the $(k+1)$-th kind* $D_{n,k}(x,a) \in \mathbb{F}_q[x]$ for $n \geq 1$ by

$$D_{n,k}(x,a) := \sum_{i=0}^{[\frac{n}{2}]} \frac{n-ki}{n-i}\binom{n-i}{i}(-a)^i x^{n-2i}$$

and $D_{0,k}(x,a) := 2-k$.

Hou, Mullen, Sellers and Yucas [4] introduced the definition of *the reversed Dickson polynomial of the first kind*, denoted by $D_n(a,x)$, as follows

Date: April 18, 2016.
2000 *Mathematics Subject Classification.* Primary 11T06, 11T55, 11C08.
Key words and phrases. Permutation polynomial, Reversed Dickson polynomial of the fourth kind, Finite field, Generating function.
*Hong is the corresponding author and was supported partially by National Science Foundation of China Grant # 11371260. Cheng was supported partially by the General Project of Department of Education of Sichuan Province # 15ZB0434. Qin was supported partially by Science and Technology Research Projects of Chongqing Education Committee Grant # KJ15012004.
Emails: ckm20@126.com, cheng.km@stu.scu.edu.cn (K. Cheng); sfhong@scu.edu.cn, s-f.hong@tom.com, hongsf02@yahoo.com (S. Hong); qincn328@sina.com (X. Qin).

$$D_n(a,x) := \sum_{i=0}^{[\frac{n}{2}]} \frac{n}{n-i}\binom{n-i}{i}(-x)^i a^{n-2i}$$

if $n \geq 1$ and $D_0(a,x) = 2$. To extend the definition of reversed Dickson polynomials, Wang and Yucas [6] defined *the n-th reversed Dickson polynomial of* $(k+1)$*-th kind* $D_{n,k}(a,x) \in \mathbb{F}_q[x]$, which is defined for $n \geq 1$ by

$$D_{n,k}(a,x) := \sum_{i=0}^{[\frac{n}{2}]} \frac{n-ki}{n-i}\binom{n-i}{i}(-x)^i a^{n-2i}$$

and $D_{0,k}(a,x) = 2-k$.

It is well known that $D_n(x,0)$ is a permutation polynomial of $\mathbb{F}_q$ if and only if $\gcd(n, q-1) = 1$, and if $a \neq 0$, then $D_n(x,a)$ induces a permutation of $\mathbb{F}_q$ if and only if $\gcd(n, q^2-1) = 1$. Besides, there are lots of published results on permutational properties of Dickson polynomial $E_n(x,a)$ of the second kind (see, for example, [1]). In [6], Wang and Yucas investigated the permutational properties of Dickson polynomial $D_{n,2}(x,1)$ of the third kind. They got some necessary conditions for $D_{n,2}(x,1)$ to be a permutation polynomial of $\mathbb{F}_q$.

Hou, Mullen, Sellers and Yucas [4] considered the permutational behavior of reversed Dickson polynomial $D_n(a,x)$ of the first kind. Actually, they showed that $D_n(a,x)$ is closely related to almost perfect nonlinear functions, and obtained some families of permutation polynomials from the revered Dickson polynomials of the first kind. In [3], Hou and Ly found several necessary conditions for the revered Dickson Polynomials $D_n(1,x)$ of the first kind to be a permutation polynomial. Recently, Hong, Qin and Zhao [2] studied the revered Dickson polynomial $E_n(a,x)$ of the second kind that is defined for $n \geq 1$ by

$$E_n(a,x) := \sum_{i=0}^{[\frac{n}{2}]} \binom{n-i}{i}(-x)^i a^{n-2i}$$

and $E_0(a,x) = 1$. In fact, they gave some necessary conditions for the revered Dickson polynomial $E_n(1,x)$ of the second kind to be a permutation polynomial of $\mathbb{F}_q$. Regarding the revered Dickson polynomial $D_{n,2}(a,x) \in \mathbb{F}_q[x]$ of the third kind, from its definition one can derive that

$$D_{n,2}(a,x) = E_{n-1}(a,x) \tag{1.1}$$

for each $x \in \mathbb{F}_q$. Using (1.1), we can deduce immediately from [2] the similar results on the permutational behavior of the reversed Dickson polynomial $D_{n,2}(a,x)$ of the third kind. Actually, for the results in [2], we need just to replace $E_n(1,x)$ by $D_{n,2}(1,x)$ and replace all other n by $n-1$, then we can obtain the corresponding results on the revered Dickson polynomial $D_{n,2}(a,x) \in \mathbb{F}_q[x]$ of the third kind. We here do not list these results.

In this paper, our main goal is to investigate the revered Dickson polynomial $D_{n,3}(a,x)$ of the fourth kind which is defined by

$$D_{n,3}(a,x) := \sum_{i=0}^{[\frac{n}{2}]} \frac{n-3i}{n-i}\binom{n-i}{i}(-x)^i a^{n-2i} \tag{1.2}$$

if $n \geq 1$ and $D_{0,3}(a,x) := -1$. For $a \neq 0$, we write $x = y(a-y)$ with an indeterminate $y \neq \frac{a}{2}$. Then $D_{n,3}(a,x)$ can be rewritten as

$$D_{n,3}(a,x)=\frac{(2a-y)y^n-(y+a)(a-y)^n}{2y-a}. \tag{1.3}$$

We have

$$D_{n,3}\Big(a,\frac{a^2}{4}\Big)=\frac{(3n-1)a^n}{2^n}. \tag{1.4}$$

In fact, (1.3) and (1.4) follows from Theorem 2.2 (i) and Theorem 2.4 (i) below. It is easy to see that $D_{n,3}(a,x)=E_n(a,x)$ if $\mathrm{char}(\mathbb{F}_q)=2$, and $D_{n,3}(a,x)=D_n(a,x)$ if $\mathrm{char}(\mathbb{F}_q)=3$. Thus we always assume $p=\mathrm{char}(\mathbb{F}_q)>3$ in what follows.

The paper is organized as follows. First in section 2, we study the properties of the reversed Dickson polynomial $D_{n,3}(a,x)$ of the fourth kind. Subsequently, in Section 3, we prove a necessary condition for the reversed Dickson polynomial $D_{n,3}(1,x)$ of the fourth kind to be a permutation polynomial of $\mathbb{F}_q$ and then introduce an auxiliary polynomial to present a characterization for $D_{n,3}(1,x)$ to be a permutation of $\mathbb{F}_q$. From the Hermite criterion [5] one knows that a function $f:\mathbb{F}_q\to\mathbb{F}_q$ is a permutation polynomial of $\mathbb{F}_q$ if and only if the i-th moment

$$\sum_{a\in\mathbb{F}_q}f(a)^i=\begin{cases}0, & \text{if } 0\le i\le q-2,\\ -1, & \text{if } i=q-1.\end{cases}$$

Thus to understand well the permutational behavior of the reversed Dickson polynomial $D_{n,3}(1,x)$ of the fourth kind, we would like to know if the i-th moment $\sum_{a\in\mathbb{F}_q}D_{n,3}(1,a)^i$ is computable. We are able to treat with this sum when $i=1$. The final section is devoted to the computation of the first moment $\sum_{a\in\mathbb{F}_q}D_{n,3}(1,a)$.

2. Revered Dickson polynomials of the fourth kind

In this section, we study the properties of the revered Dickson polynomials $D_{n,3}(a,x)$ of the fourth kind. Clearly, if $a=0$, then

$$D_{n,3}(0,x)=\begin{cases}0, & \text{if } n \text{ is odd},\\ (-1)^{\frac{n}{2}+1}x^{\frac{n}{2}}, & \text{if } n \text{ is even}.\end{cases}$$

Therefore, $D_{n,3}(0,x)$ is a PP (permutation polynomial) of $\mathbb{F}_q$ if and only if n is an even integer with $\gcd(\frac{n}{2},q-1)=1$. In what follows, we always let $a\in\mathbb{F}_q^*$. First, we give a basic fact as follows.

Lemma 2.1. [5] *Let $f(x)\in\mathbb{F}_q[x]$. Then $f(x)$ is a PP of $\mathbb{F}_q$ if and only if $cf(dx)$ is a PP of $\mathbb{F}_q$ for any given $c,d\in\mathbb{F}_q^*$.*

Then we can deduce the following result.

Theorem 2.2. *Let $a,b\in\mathbb{F}_q^*$. Then the following are true.*

(i). *One has $D_{n,3}(a,x)=\frac{a^n}{b^n}D_{n,3}(b,\frac{b^2}{a^2}x)$.*

(ii). *We have that $D_{n,3}(a,x)$ is a PP of $\mathbb{F}_q$ if and only if $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$.*

Proof. (i). By the definition of $D_{n,3}(a,x)$, we have

$$\begin{aligned}
&\frac{a^n}{b^n}D_{n,3}\Big(b,\frac{b^2}{a^2}x\Big)\\
=&\frac{a^n}{b^n}\sum_{i=0}^{[\frac{n}{2}]}\frac{n-3i}{n-i}\binom{n-i}{i}(-1)^i b^{n-2i}\frac{b^{2i}}{a^{2i}}x^i\\
=&\sum_{i=0}^{[\frac{n}{2}]}\frac{n-3i}{n-i}\binom{n-i}{i}(-1)^i a^{n-2i}x^i\\
=&D_{n,3}(a,x)
\end{aligned}$$

as required. Part (i) is proved.

(ii). Taking $b=1$ in part (i), we have

$$D_{n,3}(a,x)=a^nD_{n,3}\Big(1,\frac{x}{a^2}\Big).$$

It then follows from Lemma 2.1 that $D_{n,3}(a,x)$ is a PP of $\mathbb{F}_q$ if and only if $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$. This completes the proof of part (ii). So Theorem 2.2 is proved. □

Theorem 2.2 tells us that to study the permutational behavior of $D_{n,3}(a,x)$ over $\mathbb{F}_q$, one only needs to consider that of $D_{n,3}(1,x)$. In the following, we supply several basic properties on the revered Dickson polynomial $D_{n,3}(1,x)$ of the fourth kind. The following result is given in [2] and [4] without proof. For the completeness, we here present a proof.

Lemma 2.3. [2] [4] *Let $n\geq 0$ be an integer. Then we have $D_n(1,x(1-x))=x^n+(1-x)^n$ and $E_n(1,x(1-x))=\frac{x^{n+1}-(1-x)^{n+1}}{2x-1}$.*

Proof. Since $D_0(1,x(1-x))=2$, the first formula is true for the case $n=0$. Let now $n\geq 1$ be an integer. Then

$$D_n(1,x(1-x))=\sum_{i=0}^{[\frac{n}{2}]}\frac{n}{n-i}\binom{n-i}{i}(x+1-x)^{n-2i}(-x(1-x))^i.$$

It then follows from Waring's formula (see, for instance, Theorem 1.76 of [5]) that for any integer $n\geq 1$, we have

$$D_n(1,x(1-x))=x^n+(1-x)^n. \tag{2.1}$$

as desired. The first formula is proved.

Since $E_0(1,x(1-x))=E_1(1,x(1-x))=1$, the second formula holds when $n=0$ and 1. Now let $n\geq 2$ be an integer. Then we have

$$
\begin{aligned}
&E_n(1,x(1-x))\\
=&\sum_{i=0}^{[\frac{n}{2}]}\frac{n-i}{n-i}\binom{n-i}{i}(-x(1-x))^i\\
=&\sum_{i=0}^{[\frac{n}{2}]}\frac{n}{n-i}\binom{n-i}{i}(-x(1-x))^i+x(1-x)\sum_{i=1}^{[\frac{n}{2}]}\frac{i}{n-i}\binom{n-i}{i}(-x(1-x))^{i-1}\\
=&\sum_{i=0}^{[\frac{n}{2}]}\frac{n}{n-i}\binom{n-i}{i}(-x(1-x))^i+x\sum_{i=1}^{[\frac{n}{2}]}\binom{n-1-i}{i-1}(-x(1-x))^{i-1}\\
=&\sum_{i=0}^{[\frac{n}{2}]}\frac{n}{n-i}\binom{n-i}{i}(-x(1-x))^i+x\sum_{i=0}^{[\frac{n-2}{2}]}\binom{n-2-i}{i}(-x(1-x))^i\\
=&D_n(1,x(1-x))+xE_{n-2}(1,x(1-x)).
\end{aligned}
$$

It follows that

$$
\begin{aligned}
&E_n(1,x(1-x))\\
=&\sum_{i=0}^{[\frac{n}{2}]-1}x^i(1-x)^iD_{n-2i}(1,x(1-x))+x^{[\frac{n}{2}]}(1-x)^{[\frac{n}{2}]}E_{n-2[\frac{n}{2}]}(1,x(1-x)).
\end{aligned}\tag{2.2}
$$

From (2.1) and (2.2) one can deduce that if $n\geq 1$ is odd, then we have

$$
\begin{aligned}
&E_n(1,x(1-x))\\
=&\sum_{i=0}^{\frac{n-3}{2}}x^i(1-x)^iD_{n-2i}(1,x(1-x))+y^{\frac{n-1}{2}}(1-x)^{\frac{n-1}{2}}E_1(1,x(1-x))\\
=&\sum_{i=0}^{\frac{n-3}{2}}x^i(1-x)^i\big(x^{n-2i}+(1-x)^{n-2i}\big)+x^{\frac{n-1}{2}}(1-x)^{\frac{n-1}{2}}(x+1-x)\\
=&\sum_{i=0}^{\frac{n-1}{2}}\big(x^{n-i}(1-x)^i+x^i(1-x)^{n-i}\big)\\
=&\sum_{i=0}^{n}x^{n-i}(1-x)^i\\
=&\frac{x^{n+1}-(1-x)^{n+1}}{2x-1},
\end{aligned}
$$

and if $n\geq 0$ is even, then one has

$$\begin{aligned}
&E_n(1, x(1-x)) \\
=&\sum_{i=0}^{\frac{n}{2}-1} x^i(1-x)^i D_{n-2i}(1, x(1-x)) + x^{\frac{n}{2}}(1-x)^{\frac{n}{2}} E_0(1, x(1-x)) \\
=&\sum_{i=0}^{\frac{n}{2}-1} \left(x^{n-i}(1-x)^i + x^i(1-x)^{n-i}\right) + x^{\frac{n}{2}}(1-x)^{\frac{n}{2}} \\
=&\sum_{i=0}^{n} x^{n-i}(1-x)^i \\
=&\frac{x^{n+1} - (1-x)^{n+1}}{2x-1}
\end{aligned}$$

as expected. So the second formula is proved.

This concludes the proof of Lemma 2.3. □

Theorem 2.4. *Each of the following is true.*

(i). *For any integer $n \geq 0$, we have $D_{n,3}(1, \frac{1}{4}) = \frac{3n-1}{2^n}$ and $D_{n,3}(1, x(1-x)) = \frac{(2-x)x^n - (x+1)(1-x)^n}{2x-1}$ if $x \neq \frac{1}{2}$.*

(ii). *If n_1 and n_2 are positive integers such that $n_1 \equiv n_2 \pmod{q^2-1}$, then one has $D_{n_1,3}(1, x_0) = D_{n_2,3}(1, x_0)$ for any $x_0 \in \mathbb{F}_q \setminus \{\frac{1}{4}\}$.*

Proof. (i). First of all, it is easy to see that $D_{0,3}\big(1, \frac{1}{4}\big) = -1 = \frac{3\times 0-1}{2^0}$ and $D_{1,3}\big(1, \frac{1}{4}\big) = 1 = \frac{3\times 1-1}{2^1}$. the first identity is true for the cases that $n = 0$ and 1. Now let $n \geq 2$. Then one has

$$\begin{aligned}
D_{n,3}\Big(1, \frac{1}{4}\Big) &= \sum_{i=0}^{[\frac{n}{2}]} \frac{n-3i}{n-i} \binom{n-i}{i} \Big(-\frac{1}{4}\Big)^i \\
&= \sum_{i=0}^{[\frac{n}{2}]} \frac{n-2i}{n-i} \binom{n-i}{i} \Big(-\frac{1}{4}\Big)^i + \sum_{i=0}^{[\frac{n}{2}]} \frac{-i}{n-i} \binom{n-i}{i} \Big(-\frac{1}{4}\Big)^i \\
&= D_{n,2}\Big(1, \frac{1}{4}\Big) + \frac{1}{4} \sum_{i=0}^{[\frac{n}{2}]-1} \binom{n-2-i}{i} \Big(-\frac{1}{4}\Big)^i \\
&= D_{n,2}\Big(1, \frac{1}{4}\Big) + \frac{1}{4} E_{n-2}\Big(1, \frac{1}{4}\Big).
\end{aligned}$$

But (1.1) gives us that $D_{n,2}(1, \frac{1}{4}) = E_{n-1}(1, \frac{1}{4})$. Hence Theorem 2.2 of [2] implies that

$$\begin{aligned}
D_{n,3}\Big(1, \frac{1}{4}\Big) =& E_{n-1}\Big(1, \frac{1}{4}\Big) + \frac{1}{4} E_{n-2}\Big(1, \frac{1}{4}\Big) \\
=& \frac{n}{2^{n-1}} + \frac{1}{4} \cdot \frac{n-1}{2^{n-2}} \\
=& \frac{3n-1}{2^n}
\end{aligned}$$

as desired. So the first identity is proved.

Now we turn our attention to the second identity. Let $x \neq \frac{1}{2}$. Then by the definition of the n-th reversed Dickson polynomial of the fourth kind, one has

$$\begin{aligned} D_{n,3}(1, x(1-x)) &= \sum_{i=0}^{[\frac{n}{2}]} \frac{n-3i}{n-i}\binom{n-i}{i}(-x(1-x))^i \\ &= \sum_{i=0}^{[\frac{n}{2}]} \frac{3(n-i)-2n}{n-i}\binom{n-i}{i}(-x(1-x))^i \\ &= 3\sum_{i=0}^{[\frac{n}{2}]} \binom{n-i}{i}(-x(1-x))^i - 2\sum_{i=0}^{[\frac{n}{2}]} \frac{n}{n-i}\binom{n-i}{i}(-x(1-x))^i \\ &= 3E_n(1, x(1-x)) - 2D_n(1, x(1-x)). \end{aligned} \tag{2.3}$$

But Lemma 2.3 gives us that

$$D_n(1, x(1-x)) = x^n + (1-x)^n \tag{2.4}$$

and

$$E_n(1, x(1-x)) = \sum_{i=0}^{n} x^{n-i}(1-x)^i = \frac{x^{n+1} - (1-x)^{n+1}}{2x-1}. \tag{2.5}$$

Thus it follows from (2.3) to (2.5) that

$$\begin{aligned} D_{n,3}(1, x) =& D_{n,3}(1, y(1-y)) \\ =& 3E_n(1, y(1-y)) - 2D_n(1, y(1-y)) \\ =& \frac{3y^{n+1} - 3(1-y)^{n+1}}{2y-1} - 2\big(y^n + (1-y)^n\big) \\ =& \frac{(2-y)y^n - (y+1)(1-y)^n}{2y-1} \end{aligned}$$

as required. So the second identity holds. Part (i) is proved.

(ii). For each $x_0 \in \mathbb{F}_q \setminus \{\frac{1}{4}\}$, one can choose an element $y_0 \in \mathbb{F}_{q^2} \setminus \{\frac{1}{2}\}$ such that $x_0 = y_0(1-y_0)$. Since $n_1 \equiv n_2 \pmod{q^2-1}$, one has $y_0^{n_1} = y_0^{n_2}$ and $(1-y_0)^{n_1} = (1-y_0)^{n_2}$. It then follows from part (i) that

$$\begin{aligned} D_{n_1,3}(1, x_0) &= D_{n_1,3}(1, y_0(1-y_0)) \\ &= \frac{(2-y_0)y_0^{n_1} - (y_0+1)(1-y_0)^{n_1}}{2y_0-1} \\ &= \frac{(2-y_0)y_0^{n_2} - (y_0+1)(1-y_0)^{n_2}}{2y_0-1} \\ &= D_{n_2,3}(1, x_0) \end{aligned}$$

as desired. This ends the proof of Theorem 2.4. □

Evidently, by Theorem 2.2 (i) and Theorem 2.4 (i) one can derive that (1.3) and (1.4) are true.

Proposition 2.5. *Let $n \geq 2$ be an integer. Then the recursion*

$$D_{n,3}(1, x) = D_{n-1,3}(1, x) - xD_{n-2,3}(1, x)$$

holds for any $x \in \mathbb{F}_q$.

Proof. We consider the following two cases.

CASE 1. $x \neq \frac{1}{4}$. For this case, one may let $x = y(1-y)$ with $y \in \mathbb{F}_{q^2} \setminus \{\frac{1}{2}\}$. Then by Theorem 2.4 (i), we have

$$\begin{aligned}
&D_{n-1,3}(1,x) - xD_{n-2,3}(1,x)\\
&= D_{n-1,3}(1, y(1-y)) - y(1-y)D_{n-2,3}(1, y(1-y))\\
&= \frac{(2-y)y^{n-1} - (y+1)(1-y)^{n-1}}{2y-1} - y(1-y)\frac{(2-y)y^{n-2} - (y+1)(1-y)^{n-2}}{2y-1}\\
&= \frac{(2-y)y^{n} - (y+1)(1-y)^{n}}{2y-1}\\
&= D_{n,3}(1,x)
\end{aligned}$$

as required.

CASE 2. $x = \frac{1}{4}$. Then by Theorem 2.4 (i), we have

$$\begin{aligned}
&D_{n-1,3}\Big(1,\frac{1}{4}\Big) - \frac{1}{4}D_{n-2,3}\Big(1,\frac{1}{4}\Big)\\
=&\frac{3n-4}{2^{n-1}} - \frac{1}{4}\frac{3n-7}{2^{n-2}}\\
=&\frac{3n-1}{2^n}\\
=&D_{n,3}\Big(1,\frac{1}{4}\Big).
\end{aligned}$$

This concludes the proof of Proposition 2.5. □

By Proposition 2.5, we can obtain the generating function of the revered Dickson polynomial $D_{n,3}(1,x)$ of the fourth kind as follows.

Proposition 2.6. *The generating function of $D_{n,3}(1,x)$ is given by*

$$\sum_{n=0}^{\infty} D_{n,3}(1,x)t^n = \frac{2t-1}{1-t+xt^2}.$$

Proof. By the recursion presented in Proposition 2.5, we have

$$\begin{aligned}
&(1-t+xt^2)\sum_{n=0}^{\infty} D_{n,3}(1,x)t^n\\
=&\sum_{n=0}^{\infty} D_{n,3}(1,x)t^n - \sum_{n=0}^{\infty} D_{n,3}(1,x)t^{n+1} + x\sum_{n=0}^{\infty} D_{n,3}(1,x)t^{n+2}\\
=&2t-1+\sum_{n=0}^{\infty}\big(D_{n+2,3}(1,x) - D_{n+1,3}(1,x) + xD_{n,3}(1,x)\big)t^{n+2}\\
=&2t-1.
\end{aligned}$$

Thus the desired result follows immediately. □

Now we can use Theorem 2.4 to present an explicit formula for $D_{n,3}(1,x)$ when n is a power of the characteristic p. Then we show that $D_{n,3}(1,x)$ is not a PP of $\mathbb{F}_q$ in this case.

Proposition 2.7. *Let $p = \mathrm{char}(\mathbb{F}_q) > 3$ and k be a positive integer. Then*

$$2^{p^k} D_{p^k,3}(1,x) + 1 = 3(1-4x)^{\frac{p^k-1}{2}}.$$

Proof. Putting $x = y(1-y)$ in Theorem 2.4 (i) gives us that

$$\begin{aligned} D_{p^k,3}(1,x) =& D_{p^k,3}(1, y(1-y)) \\ =& \frac{(2-y)y^{p^k} - (y+1)(1-y)^{p^k}}{2y-1} \\ =& \frac{\frac{3-u}{2}\left(\frac{u+1}{2}\right)^{p^k} - \frac{3+u}{2}\left(\frac{1-u}{2}\right)^{p^k}}{u} \\ =& \frac{1}{2^{p^k+1}u}\Big((3-u)(u+1)^{p^k} - (u+3)(1-u)^{p^k}\Big) \\ =& \frac{1}{2^{p^k}}(3u^{p^k-1} - 1), \end{aligned}$$

where $u = 2y - 1$. So we obtain that

$$\begin{aligned} & 2^{p^k} D_{p^k,3}(1,x) \\ =& 3(u^2)^{\frac{p^k-1}{2}} - 1 \\ =& 3\big((2y-1)^2\big)^{\frac{p^k-1}{2}} - 1, \end{aligned}$$

which infers that

$$2^{p^k} D_{p^k,3}(1,x) + 1 = 3(1-4x)^{\frac{p^k-1}{2}}$$

as desired. So Proposition 2.7 is proved. □

It is well known that every linear polynomial over $\mathbb{F}_q$ is a PP of $\mathbb{F}_q$ and that the monomial x^n is a PP of $\mathbb{F}_q$ if and only if $\gcd(n, q-1) = 1$. Then by Proposition 2.7, we have the following result.

Corollary 2.8. *Let $p > 3$ be a prime and $q = p^e$. Let e and k be positive integers with $k \le e$. Then $D_{p^k,3}(1,x)$ is not a PP of $\mathbb{F}_q$.*

Proof. By Proposition 2.7, we know that $D_{p^k,3}(1,x)$ is a PP of $\mathbb{F}_q$ if and only if

$$(1-4x)^{\frac{p^k-1}{2}}$$

is a PP of $\mathbb{F}_q$ which is equivalent to

$$\gcd\Big(\frac{p^k-1}{2}, q-1\Big) = 1.$$

The latter one is impossible since $\frac{p-1}{2} \mid \gcd\left(\frac{p^k-1}{2}, q-1\right)$ implies that

$$\gcd\Big(\frac{p^k-1}{2}, q-1\Big) \ge \frac{p-1}{2} > 1.$$

Thus $D_{p^k,3}(1,x)$ is not a PP of $\mathbb{F}_q$. □

Lemma 2.9. [4] *Let $x \in \mathbb{F}_{q^2}$. Then $x(1-x) \in \mathbb{F}_q$ if and only if $x^q = x$ or $x^q = 1 - x$.*

Let V be defined by

$$V := \{x \in \mathbb{F}_{q^2} : x^q = 1 - x\}.$$

Clearly, $\mathbb{F}_q \cap V = \{\frac{1}{2}\}$. Then we obtain a characterization for $D_{n,3}(1,x)$ to be a PP of $\mathbb{F}_q$ as follows.

Theorem 2.10. *Let $q = p^e$ with $p > 3$ being a prime and e being a positive integer. Let*

$$f : y \mapsto \frac{(2-y)y^n - (y+1)(1-y)^n}{2y-1}$$

be a mapping on $(\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$. Then $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$ if and only if f is 2-to-1 and $f(y) \neq \frac{3n-1}{2^n}$ for any $y \in (\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$.

Proof. First, we show the sufficiency part. Let f be 2-to-1 and $f(y) \neq \frac{3n-1}{2^n}$ for any $y \in (\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$. Let $D_{n,3}(1,x_1) = D_{n,3}(1,x_2)$ for $x_1, x_2 \in \mathbb{F}_q$. To show that $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$, it suffices to show that $x_1 = x_2$ that will be done in what follows.

First of all, one can find $y_1, y_2 \in \mathbb{F}_{q^2}$ satisfying $x_1 = y_1(1-y_1)$ and $x_2 = y_2(1-y_2)$. By Lemma 2.9, we know that $y_1, y_2 \in \mathbb{F}_q \cup V$. We divide the proof into the following two cases.

Case 1. At least one of x_1 and x_2 is equal to $\frac{1}{4}$. Without loss of any generality, we may let $x_1 = \frac{1}{4}$. So by Theorem 2.4 (i), one derives that

$$D_{n,3}(1,x_2) = D_{n,3}(1,x_1) = D_{n,3}\Big(1,\frac{1}{4}\Big) = \frac{3n-1}{2^n}. \tag{2.6}$$

We claim that $x_2 = \frac{1}{4}$. Assume that $x_2 \neq \frac{1}{4}$. Then $y_2 \neq \frac{1}{2}$. Since $f(y) \neq \frac{3n-1}{2^n}$ for any $y \in (\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$, by Theorem 2.4 (i), we get that

$$D_{n,3}(1,x_2) = \frac{(2-y_2)y_2^n - (y_2+1)(1-y_2)^n}{2y_2-1} = f(y_2) \neq \frac{3n-1}{2^n},$$

which contradicts to (2.6). Hence the claim is true, and so we have $x_1 = x_2$ as required.

Case 2. Both of x_1 and x_2 are not equal to $\frac{1}{4}$. Then $y_1 \neq \frac{1}{2}$ and $y_2 \neq \frac{1}{2}$. Since $D_{n,3}(1,x_1) = D_{n,3}(1,x_2)$, by Theorem 2.4 (i), one has

$$\frac{(2-y_2)y_1^n - (y_1+1)(1-y_1)^n}{2y_1-1} = \frac{(2-y_2)y_2^n - (y_2+1)(1-y_2)^n}{2y_2-1},$$

which is equivalent to $f(y_1) = f(y_2)$. However, f is a 2-to-1 mapping on $(\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$, and $f(y_2) = f(1-y_2)$ by the definition of f. It then follows that $y_1 = y_2$ or $y_1 = 1-y_2$. Thus $x_1 = x_2$ as desired. Hence the sufficiency part is proved.

Now we prove the necessity part. Let $D_{n,3}(1,x)$ be a PP of $\mathbb{F}_q$. Choose two elements $y_1, y_2 \in (\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$ such that $f(y_1) = f(y_2)$, that is,

$$\frac{(2-y_1)y_1^n - (y_1+1)(1-y_1)^n}{2y_1-1} = \frac{(2-y_2)y_2^n - (y_2+1)(1-y_2)^n}{2y_2-1}. \tag{2.7}$$

Since $y_1, y_2 \in (\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$, it follows from Lemma 2.9 that $y_1(1-y_1) \in \mathbb{F}_q$ and $y_2(1-y_2) \in \mathbb{F}_q$. So by Theorem 2.4 (i), (2.7) implies that

$$D_{n,3}(1, y_0(1-y_0)) = D_{n,3}(1, y(1-y)).$$

Thus $y_1(1-y_1) = y_2(1-y_2)$ since $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$, which infers that $y_1 = y_2$ or $y_1 = 1-y_2$. Since $y_2 \neq \frac{1}{2}$, one has $y_2 \neq 1-y_2$. Therefore f is a 2-to-1 mapping on $(\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$.

Now take $y' \in (\mathbb{F}_q \cup V) \setminus \{\frac{1}{2}\}$. Then from Lemma 2.9 it follows that $y'(1-y') \in \mathbb{F}_q$ and

$$y'(1-y') \neq \frac{1}{2}\Big(1-\frac{1}{2}\Big).$$

Notice that $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$. Hence one has

$$D_{n,3}(1, y'(1-y')) \neq D_{n,3}\Big(1, \frac{1}{2}\Big(1-\frac{1}{2}\Big)\Big).$$

But Theorem 2.4 (i) tells us that

$$D_{n,3}\Big(1,\frac{1}{2}\Big(1-\frac{1}{2}\Big)\Big)=\frac{3n-1}{2^n}.$$

Then by Theorem 2.4 (i) and noting that $y'\neq\frac{1}{2}$, we have

$$\frac{(2-y')y'^n-(y'+1)(1-y')^n}{2y'-1}\neq\frac{3n-1}{2^n},$$

which infers that $f(y')\neq\frac{3n-1}{2^n}$ for any $y'\in(\mathbb{F}_q\cup V)\setminus\{\frac{1}{2}\}$. So the necessity part is proved.

The proof of Theorem 2.10 is complete. □

3. A necessary condition for $D_{n,3}(1,x)$ to be permutational and an auxiliary polynomial

In this section, we study some necessary conditions on n for $D_{n,3}(1,x)$ to be a PP of $\mathbb{F}_q$. It is easy to check that

$$D_{n,3}(1,0)=1, D_{0,3}(1,1)=-1, D_{1,3}(1,1)=1, D_{0,3}(1,-2)=-1, D_{1,3}(1,-2)=1.$$

Then by Proposition 2.5, we have the following recursion relations

$$\begin{cases} D_{0,3}(1,1)=-1,\\ D_{1,3}(1,1)=1,\\ D_{n+2,3}(1,1)=D_{n+1,3}(1,1)-D_{n,3}(1,1),\end{cases}$$

and

$$\begin{cases} D_{0,3}(1,-2)=-1,\\ D_{1,3}(1,-2)=1,\\ D_{n+2,3}(1,-2)=D_{n+1,3}(1,-2)+2D_{n,3}(1,-2).\end{cases}$$

From these recursive formulas, one can easily show that the sequences

$$\{D_{n,3}(1,1)|n\in\mathbb{N}\} \text{ and } \{D_{n,3}(1,-2)|n\in\mathbb{N}\}$$

are periodic with the smallest periods 6 and 2, respectively. In fact, one has

$$D_{n,3}(1,1)=\begin{cases} 1, & \text{if } n\equiv 1,3 \pmod 6,\\ -1, & \text{if } n\equiv 0,4 \pmod 6,\\ 2, & \text{if } n\equiv 2 \pmod 6,\\ -2, & \text{if } n\equiv 5 \pmod 6\end{cases}$$

and

$$D_{n,3}(1,-2)=\begin{cases} 1, & \text{if } n\equiv 1 \pmod 2,\\ -1, & \text{if } n\equiv 0 \pmod 2.\end{cases}$$

Theorem 3.1. *Assume that $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$ with $q=p^e$ and $p>3$. Then $n\equiv 2 \pmod 6$.*

Proof. Let $D_{n,3}(1,x)$ be a PP of $\mathbb{F}_q$. Then $D_{n,3}(1,0), D_{n,3}(1,1)$ and $D_{n,3}(1,-2)$ are distinct. Since $D_{n,3}(1,0)=1$, one has $D_{n,3}(1,1)\neq 1$ and $D_{n,3}(1,-2)\neq 1$. Then the above results tells us that $n\not\equiv 1,3,5 \pmod 6$. Further, we have $D_{n,3}(1,-2)=-1$ which means that n must be even, and so $D_{n,3}(1,1)\neq -2$. But $D_{n,3}(1,1)\neq D_{n,3}(1,-2)$. So $D_{n,3}(1,1)\neq -1$. Hence $D_{n,3}(1,1)=2$. Finally, the desired result $n\equiv 2 \pmod 6$ follows immediately. □

Evidently, Corollary 2.8 can be easily deduced from Theorem 3.1. Furthermore, By Theorem 3.1, we know that $D_{n,3}(1,x)$ is not a PP of $\mathbb{F}_q$ if n is odd.

In what follows, we investigate $D_{n,3}(1,x)$ with n being an even number. We define the following auxiliary polynomial $f_n(x) \in \mathbb{Z}[x]$ by

$$f_n(x) := -x^{\frac{n}{2}} + \sum_{j=0}^{\frac{n}{2}-1} \frac{3n-8j-1}{n+1}\binom{n+1}{2j+1}x^j.$$

Then we have the following relation between $D_{n,3}(1,x)$ and $f_n(x)$.

Theorem 3.2. *Let $p>3$ be a prime and $n \geq 0$ be an even integer. Then*

(i). *One has*

$$D_{n,3}(1,x) = \frac{1}{2^n} f_n(1-4x). \tag{3.1}$$

(ii). *We have that $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$ if and only if $f_n(x)$ is a PP of $\mathbb{F}_q$.*

Proof. (i). First, let $x \in \mathbb{F}_q \setminus \{\frac{1}{4}\}$. Then there exists $y \in \mathbb{F}_{q^2} \setminus \{\frac{1}{2}\}$ such that $x = y(1-y)$. Let $u = 2y-1$. Since for any integer j with $0 \leq j \leq \frac{n}{2}-1$, one has

$$3\binom{n}{2j+1} - \binom{n}{2j} = \frac{3n-8j-1}{n+1}\binom{n+1}{2j+1},$$

it then follows from Theorem 2.4 (i) that

$$\begin{aligned}
D_{n,3}(1,x) &= D_{n,3}(1, y(1-y)) \\
&= \frac{(2-y)y^n - (y+1)(1-y)^n}{2y-1} \\
&= \frac{\frac{3-u}{2}\left(\frac{u+1}{2}\right)^n - \frac{3+u}{2}\left(\frac{1-u}{2}\right)^n}{u} \\
&= \frac{1}{2^{n+1}u}\big((3-u)(u+1)^n - (u+3)(1-u)^n\big) \\
&= \frac{1}{2^n}\Big(-u^n + \sum_{j=0}^{\frac{n}{2}-1}\Big(3\binom{n}{2j+1} - \binom{n}{2j}\Big)u^{2j}\Big) \\
&= \frac{1}{2^n} f_n(u^2) \\
&= \frac{1}{2^n} f_n(1-4y(1-y)) \\
&= \frac{1}{2^n} f_n(1-4x)
\end{aligned}$$

as desired. So (3.1) holds in this case.

Consequently, we let $x = \frac{1}{4}$. Then by Theorem 2.4 (i), we have

$$D_{n,3}\Big(1, \frac{1}{4}\Big) = \frac{3n-1}{2^n}.$$

On the other hand, we can easily check that $f_n(0) = 3n-1$. Therefore

$$D_{n,3}\Big(1, \frac{1}{4}\Big) = \frac{1}{2^n} f_n(0) = \frac{1}{2^n} f_n\Big(1 - 4 \times \frac{1}{4}\Big)$$

as one desires. So (3.1) is proved.

(ii). Notice that $\frac{1}{2^n} \in \mathbb{F}_q^*$ and $1-4x$ is linear. So $D_{n,3}(1,x)$ is a PP of $\mathbb{F}_q$ if and only if $f_n(x)$ is a PP of $\mathbb{F}_q$. This ends the proof of Theorem 3.2. □

4. The first moment $\sum_{a\in\mathbb{F}_q} D_{n,3}(1,a)$

In this section, we compute the first moment $\sum_{a\in\mathbb{F}_q} D_{n,3}(1,a)$. By Proposition 2.6, one has

$$\begin{aligned}\sum_{n=0}^{\infty} D_{n,3}(1,x)t^n &= \frac{2t-1}{1-t+xt^2} = \frac{2t-1}{1-t}\frac{1}{1-\frac{t^2}{t-1}x}\\ &= \frac{2t-1}{1-t}\Big(1+\sum_{k=1}^{q-1}\sum_{\ell=0}^{\infty}\Big(\frac{t^2}{t-1}\Big)^{k+\ell(q-1)}x^{k+\ell(q-1)}\Big)\\ &\equiv \frac{2t-1}{1-t}\Big(1+\sum_{k=1}^{q-1}\sum_{\ell=0}^{\infty}\Big(\frac{t^2}{t-1}\Big)^{k+\ell(q-1)}x^{k}\Big) \pmod{x^q-x}\\ &= \frac{2t-1}{1-t}\Big(1+\sum_{k=1}^{q-1}\frac{(\frac{t^2}{t-1})^k}{1-(\frac{t^2}{t-1})^{q-1}}x^k\Big)\\ &= \frac{2t-1}{1-t}\Big(1+\sum_{k=1}^{q-1}\frac{(t-1)^{q-1-k}t^{2k}}{(t-1)^{q-1}-t^{2(q-1)}}x^k\Big). \end{aligned} \tag{4.1}$$

Moreover, by Theorem 2.4 (ii), it follows that for any $x\in\mathbb{F}_q\setminus\{\frac{1}{4}\}$, one has

$$D_{n_1,3}(1,x) = D_{n_2,3}(1,x)$$

when $n_1\equiv n_2 \pmod{q^2-1}$. Thus if $x\neq\frac{1}{4}$, one has

$$\begin{aligned}\sum_{n=0}^{\infty} D_{n,3}(1,x)t^n &= 1+\sum_{n=1}^{q^2-1}\sum_{\ell=0}^{\infty} D_{n+\ell(q^2-1),3}(1,x)t^{n+\ell(q^2-1)}\\ &= 1+\sum_{n=1}^{q^2-1} D_{n,3}(1,x)\sum_{\ell=0}^{\infty} t^{n+\ell(q^2-1)}\\ &= 1+\frac{1}{1-t^{q^2-1}}\sum_{n=1}^{q^2-1} D_{n,3}(1,x)t^n. \end{aligned} \tag{4.2}$$

Then (4.1) together with (4.2) gives that for any $x\neq\frac{1}{4}$, we have

$$\begin{aligned}&\sum_{n=1}^{q^2-1} D_{n,3}(1,x)t^n \qquad (4.3)\\ =&\Big(\sum_{n=0}^{\infty} D_{n,3}(1,x)t^n - 1\Big)(1-t^{q^2-1})\\ \equiv&\Big(\frac{2t-1}{1-t}-1\Big)(1-t^{q^2-1}) + \frac{(1-t^{q^2-1})(2t-1)}{1-t}\sum_{k=1}^{q-1}\frac{(t-1)^{q-1-k}t^{2k}}{(t-1)^{q-1}-t^{2(q-1)}}x^k \pmod{x^q-x}\\ =&\frac{(3t-2)(1-t^{q^2-1})}{1-t} + h(t)\sum_{k=1}^{q-1}(t-1)^{q-1-k}t^{2k}x^k, \qquad (4.4)\end{aligned}$$

where

$$h(t) := \frac{(t^{q^2-1}-1)(2t-1)}{(t-1)^q-(t-1)t^{2(q-1)}}.$$

Lemma 4.1. [5] *Let $u_0, u_1, \cdots, u_{q-1}$ be the list of the all elements of $\mathbb{F}_q$. Then*

$$\sum_{i=0}^{q-1} u_i^k = \begin{cases} 0, & \text{if } 0 \le k \le q-2, \\ -1, & \text{if } k = q-1. \end{cases}$$

Now by Theorem 2.4 (i), Lemma 4.1 and (4.4), we derive that

$$\begin{aligned}
&\sum_{n=1}^{q^2-1} \sum_{a\in\mathbb{F}_q} D_{n,3}(1,a)t^n \\
=&\sum_{n=1}^{q^2-1} D_{n,3}\Big(1,\frac{1}{4}\Big)t^n + \sum_{n=1}^{q^2-1} \sum_{a\in\mathbb{F}_q\setminus\{\frac{1}{4}\}} D_{n,3}(1,a)t^n \\
=&\sum_{n=1}^{q^2-1} \frac{3n-1}{2^n}t^n + \sum_{a\in\mathbb{F}_q\setminus\{\frac{1}{4}\}} \frac{(3t-2)(1-t^{q^2-1})}{1-t} + h(t)\sum_{k=1}^{q-1}(t-1)^{q-1-k}t^{2k} \sum_{a\in\mathbb{F}_q\setminus\{\frac{1}{4}\}} a^k \\
=&\sum_{n=1}^{q^2-1} \frac{3n-1}{2^n}t^n + (q-1)\frac{(3t-2)(1-t^{q^2-1})}{1-t} + h(t)\sum_{k=1}^{q-1}(t-1)^{q-1-k}t^{2k} \sum_{a\in\mathbb{F}_q} a^k \\
&- h(t)\sum_{k=1}^{q-1}(t-1)^{q-1-k}t^{2k}\Big(\frac{1}{4}\Big)^k \\
=&\sum_{n=1}^{q^2-1} \frac{3n-1}{2^n}t^n - \frac{(3t-2)(1-t^{q^2-1})}{1-t} - h(t)t^{2(q-1)} - h(t)\sum_{k=1}^{q-1}(t-1)^{q-1-k}t^{2k}\Big(\frac{1}{4}\Big)^k.
\end{aligned} \tag{4.5}$$

Since $(t-1)^q = t^q - 1$ and q is odd, one has

$$\begin{aligned}
h(t) =& \frac{(t^{q^2-1}-1)(2t-1)}{(t-1)^q - (t-1)t^{2(q-1)}} \\
=& \frac{(t^{q^2-1}-1)(2t-1)}{(1-t^{q-1})(t^q - t^{q-1} - 1)} \\
=& \frac{(t^{q^2}-t)(2t-1)}{(t-t^q)(t^q - t^{q-1} - 1)} \\
=& \frac{(t^q-t)^q + t^q - t}{t-t^q} \cdot \frac{2t-1}{t^q - t^{q-1} - 1} \\
=& \frac{(-1-(t-t^q)^{q-1})(2t-1)}{t^q - t^{q-1} - 1} \\
=& \frac{(2t-1)\sum_{i=0}^{q^2-q} b_i t^i}{t^q - t^{q-1} - 1},
\end{aligned} \tag{4.6}$$

where

$$\sum_{i=0}^{q^2-q} b_i t^i := -1 - (t-t^q)^{q-1}.$$

Then by the binomial theorem applied to $(t-t^q)^{q-1}$, we can derive the following expression for the coefficient b_i.

Proposition 4.2. *For each integer i with $0 \le i \le q^2-q$, write $i=\alpha+\beta q$ with α and β being integers such that $0 \le \alpha, \beta \le q-1$. Then*

$$b_i = \begin{cases} (-1)^{\beta+1}\binom{q-1}{\beta}, & \text{if } \alpha+\beta = q-1, \\ -1, & \text{if } \alpha=\beta=0, \\ 0, & \text{otherwise}. \end{cases}$$

For convenience, let

$$a_n := \sum_{a\in\mathbb{F}_q} D_{n,3}(1,a).$$

Then by (4.5) and (4.6), we arrive at

$$\sum_{n=1}^{q^2-1}\Big(a_n - \frac{3n-1}{2^n}\Big)t^n \tag{4.7}$$

$$= -\frac{(3t-2)(1-t^{q^2-1})}{1-t} - \frac{(2t-1)\sum_{i=1}^{q^2-q} b_i t^i}{t^q - t^{q-1}-1}\Big(t^{2(q-1)} + \sum_{k=1}^{q-1}(t-1)^{q-1-k}t^{2k}\Big(\frac{1}{4}\Big)^k\Big), \tag{4.8}$$

which implies that

$$(t^q - t^{q-1}-1)\sum_{n=1}^{q^2-1}\Big(a_n - \frac{3n-1}{2^n}\Big)t^n$$

$$= -(t^q - t^{q-1}-1)(3t-2)\sum_{i=0}^{q^2-2} t^i - (2t-1)\Big(t^{2(q-1)} + \sum_{k=1}^{q-1}(t-1)^{q-1-k}t^{2k}\Big(\frac{1}{4}\Big)^k\Big)\sum_{i=0}^{q^2-q} b_i t^i. \tag{4.9}$$

Let

$$\sum_{i=1}^{q^2+q-1} c_i t^i$$

denote the right-hand side of (4.9) and let

$$d_n := a_n - \frac{3n-1}{2^n}$$

for each integer n with $1 \le n \le q^2-1$. Then (4.9) can be reduced to

$$(t^q - t^{q-1}-1)\sum_{n=1}^{q^2-1} d_n t^n = \sum_{i=1}^{q^2+q-1} c_i t^i. \tag{4.10}$$

Then by comparing the coefficient of t^i with $1 \le i \le q^2+q-1$ of the both sides in (4.10), we derive the following relations:

$$\begin{cases} c_j = -d_j, & \text{if } 1 \le j \le q-1, \\ c_q = -d_1 - d_q, & \\ c_{q+j} = d_j - d_{j+1} - d_{q+j}, & \text{if } 1 \le j \le q^2-q-1, \\ c_{q^2+j} = d_{q^2-q+j} - d_{q^2-q+j+1}, & \text{if } 0 \le j \le q-2, \\ c_{q^2+q-1} = d_{q^2-1}, & \end{cases}$$

from which we can deduce that

$$\begin{cases} d_j = -c_j, & \text{if } 1 \le j \le q-1, \\ d_q = c_1 - c_q, & \\ d_{\ell q+j} = d_{(\ell-1)q+j} - d_{(\ell-1)q+j+1} - c_{\ell q+j}, & \text{if } 1 \le \ell \le q-2 \text{ and } 1 \le j \le q-1, \\ d_{\ell q} = d_{(\ell-1)q} - d_{(\ell-1)q+1} - c_{\ell q}, & \text{if } 2 \le \ell \le q-2, \\ d_{q^2-q+j} = \sum_{i=j}^{q-1} c_{q^2+i}, & \text{if } 0 \le j \le q-1. \end{cases} \tag{4.11}$$

Finally, (4.11) together with the following identity

$$\sum_{a\in\mathbb{F}_q} D_{n,3}(1,a) = d_n + \frac{3n-1}{2^n}$$

shows that the last main result of this paper is true:

Theorem 4.3. *Let c_i be the coefficient of t^i in the right-hand side of (4.9) with i being an integer such that $1 \le i \le q^2+q-1$. Then we have*

$$\sum_{a\in\mathbb{F}_q} D_{j,3}(1,a) = -c_j + \frac{3j-1}{2^j} \quad if \quad 1 \le j \le q-1,$$

$$\sum_{a\in\mathbb{F}_q} D_{q,3}(1,a) = c_1 - c_q - \frac{1}{2},$$

$$\sum_{a\in\mathbb{F}_q} D_{\ell q+j,3}(1,a) = \sum_{a\in\mathbb{F}_q} D_{(\ell-1)q+j,3}(1,a) - \sum_{a\in\mathbb{F}_q} D_{(\ell-1)q+j+1,3}(1,a) - c_{\ell q+j} + \frac{3}{2^{\ell+j}}$$
$$if \quad 1 \le \ell \le q-2 \text{ and } 1 \le j \le q-1,$$

$$\sum_{a\in\mathbb{F}_q} D_{\ell q,3}(1,a) = \sum_{a\in\mathbb{F}_q} D_{(\ell-1)q,3}(1,a) - \sum_{a\in\mathbb{F}_q} D_{(\ell-1)q+1,3}(1,a) - c_{\ell q} + \frac{3}{2^\ell} \quad if \quad 2 \le \ell \le q-2$$

and

$$\sum_{a\in\mathbb{F}_q} D_{q^2-q+j,3}(1,a) = \sum_{i=j}^{q-1} c_{q^2+i} + \frac{3j-1}{2^j} \quad if \quad 0 \le j \le q-1.$$

References

[1] S.D. Cohen, Dickson polynomials of the second kind that are permutations, Canad. J. Math. 46 (1994), 225-238.

[2] S. Hong, X. Qin and W. Zhao, Necessary conditions for reversed Dickson polynomials of the second kind to be permutational, Finite Fields Appl. 37 (2016), 54-71.

[3] X. Hou and T. Ly, Necessary conditions for reversed Dickson polynomials to be permutational, Finite Fields Appl. 16 (2010), 436-448.

[4] X. Hou, G.L. Mullen, J.A. Sellers and J.L. Yucas, Reversed Dickson polynomials over finite fields, Finite Fields Appl. 15 (2009), 748-773.

[5] R. Lidl and H. Niederreiter, Finite Fields, second ed., Encyclopedia of Mathematics and its Applications, vol.20, Cambridge University Press, Cambridge, 1997.

[6] Q. Wang and J.L. Yucas, Dickson polynomials over finite fields, Finite Fields Appl. 18 (2012), 814-831.